

Systemic Diseases and Syndromes that Affect the Periodontium

This chart includes medical conditions known to impact periodontal health and that may be included in a differential diagnosis when periodontitis is detected in pediatric patients. Individualized at-home and professional preventive oral care interventions must be emphasized for these patients. A multidisciplinary approach may be indicated for safe and effective oral health care.

Disorder	General characteristics	Diagnostic criteria	Oral findings		Treatment considerations
			Clinical	Radiographic	
Chédiak-Higashi syndrome ^{1,2}	<ul style="list-style-type: none"> - Rare autosomal recessive disorder of the immune system - Mild 'atypical' phenotype: 10-15% of cases; 'classic' phenotype: 80-90%, progresses to accelerated phase (fatal without bone marrow transplant) - Characterized by reduced pigmentation, neurological deficits, fever, lymphadenopathy, anemia, neutropenia, hepatosplenomegaly thrombocytopenia. 	<ul style="list-style-type: none"> - Microscopic analysis of WBCs - Genetic testing identifies mutations in the lysosomal trafficking regulator gene (LYST/CHS1) 	<ul style="list-style-type: none"> - Severe gingival inflammation, swelling, and recession - Early onset periodontitis in primary and permanent dentitions - Premature tooth loss - Oral ulcerations may be present 	<p>Alveolar bone loss (localized or generalized)</p>	<ul style="list-style-type: none"> - Immune status and transplantation dictate timing and precautions (e.g., antibiotics) - Supportive management of complications (e.g., antibiotics to treat bacterial infections) - Aggressive recurrent periodontitis may not respond to SRP or antibiotic treatment - For extractions/surgeries, consider adjunctive measures for hemostasis and avoid NSAIDs due to platelet dysfunction - Prosthetic therapy for lost teeth may be considered depending on patient's medical status
Diabetes mellitus ^{3,4}	<ul style="list-style-type: none"> - Metabolic disorder <ul style="list-style-type: none"> • Type 1: autoimmune reaction causes lack of insulin production; usually diagnosed in children and young adults • Type 2: insulin resistance; usually diagnosed in adults • Gestational: insulin resistance during pregnancy - Associated with increased inflammation, impaired immunologic response and wound healing, CV disease, retinopathy, nephropathy, neuropathy 	<ul style="list-style-type: none"> - Glycated hemoglobin (A1C) - Other tests include: oral glucose tolerance tests, fasting plasma glucose test, random plasma glucose test 	<ul style="list-style-type: none"> - Enlarged, erythematous attached gingiva - Dental/ periodontal abscesses - Increased clinical attachment loss and pathologic periodontal pockets - Severe periodontitis 	<p>Alveolar bone loss</p>	<ul style="list-style-type: none"> - Assess level of disease control (e.g., compliance with diet and medications) <ul style="list-style-type: none"> • With uncontrolled diabetes, consider antibiotic prophylaxis for invasive oral procedures • Nonsurgical periodontal therapy (e.g., SRP and antimicrobial agents [chlorhexidine, antibiotics]) shows modest glycemic control improvement • Monitor for delayed healing
Haim-Munk syndrome ^{4,5}	<ul style="list-style-type: none"> - Rare autosomal recessive syndrome; a phenotypic variant of PLS with mutation to chromosome 11q14-q21 and loss of function of the cathepsin C gene - Milder periodontal disease and more severe dermatologic manifestations than PLS - Clinical manifestations <ul style="list-style-type: none"> • Dermatologic: palmo-plantar hyperkeratosis; scaly patches on eyelids, lips, cheeks; skin infections • Skeletal: arachnodactyly, onychogryphosis, acroosteolysis, pes planus, muscle contractures, and destructive arthritis 	<p>Genetic testing for mutation of cathepsin C gene</p>	<ul style="list-style-type: none"> - Rapidly advancing gingival inflammation and bleeding, deep periodontal pockets, gingival abscesses, periodontal destruction - Premature loss of all primary teeth by age 4-5; loss of permanent teeth by age 16 - After tooth loss, gingiva returns to healthy state 	<p>Generalized alveolar bone loss with migration of teeth</p>	<ul style="list-style-type: none"> - Treatment of oral manifestations depends on patient's age, psychological state, and tooth mobility - May include nonsurgical therapy (e.g., monthly SRP, systemic antibiotics) and/or extraction of hopeless teeth - Alveolar loss renders prosthetic rehabilitation challenging

Table continued on the next page.

Abbreviations in table: BMI: Body mass index; CDC: Centers for Disease Control and Prevention; CV: cardiovascular; NSAIDs: nonsteroidal anti-inflammatory drugs; PLS: Papillon Lefèvre syndrome; SRP: Scaling and root planing; WBCs: White blood cells.

Disorder	General characteristics	Diagnostic criteria	Oral findings		Treatment considerations
			Clinical	Radiographic	
Hypophosphatasia^{6,7}	<ul style="list-style-type: none"> - Rare genetic metabolic bone disorder characterized by impaired mineralization of bones and/or teeth - Mutations in the ALPL gene leading to low alkaline phosphatase activity - Wide-ranging severity (involving skeletal, renal, neurological, muscular, respiratory complications); six types based on severity and age of onset - Disturbed cementum formation; tooth loss is one of the first signs of the condition 	<ul style="list-style-type: none"> - Serum alkaline phosphatase (ALP) levels - ALPL gene testing 	<ul style="list-style-type: none"> - Premature exfoliation of primary teeth with little or no root resorption - Clinical inflammation milder than in other systemic diseases associated with periodontitis - Permanent dentition may be clinically normal in mild subtypes or prone to periodontitis and early tooth loss in severe cases 	<ul style="list-style-type: none"> - Alveolar bone loss - Large pulpal chambers and root canals - Thin dentin 	<ul style="list-style-type: none"> - Enzyme replacement therapy, a new disease-modifying treatment, has shown periodontal, tooth, and bone improvements - Caution with orthodontic management due to cementum dysplasia and impaired periodontal attachment - Prosthodontic therapy may include removable prostheses and possible implants to stabilize prosthesis in the permanent dentition for those skeletally mature
Langerhans cell histiocytosis⁸	<ul style="list-style-type: none"> - Rare cancer-like condition (inflammatory myeloid neoplasia) characterized by excessive proliferation/infiltration of histiocytes (Langerhans cells); form categorized as a single-system (single organ affected) or multisystem (several organs affected) - Mortality rate: < 10% in single-system vs. 30-50% in multisystem - Average age of onset: 1-3 years; male predilection - Oral manifestations and pain can be the first signs of the condition 	<p>Clinical, microscopic, hematologic, and imaging examinations</p>	<ul style="list-style-type: none"> - Gingivitis, bleeding, recession, mucosal swelling, periodontitis, ulceration - Excessive mobility of teeth, premature exfoliation - Oral pain 	<ul style="list-style-type: none"> - Alveolar bone loss with distinct appearance of teeth floating in soft tissue - Unifocal or multiple lesions within the body of the maxilla and mandible 	<ul style="list-style-type: none"> - Management of periodontal disease is not the first-line of treatment - Treatment of oral lesions depends on the type/ extent of disease and may vary from observation to pharmacotherapy, surgical excision/curettage, and/or radiation therapy - Treatment should include basic periodontal therapy and extractions of hopeless teeth depending on immune status
Leukocyte adhesion deficiency syndromes^{4,9}	<ul style="list-style-type: none"> - Rare autosomal recessive disorders - Primary immunodeficiency disorder involving both B and T cells - Impaired migration of WBCs to infection sites - Recurrent nonpyogenic bacterial and fungal mucosal infections - Compromised wound healing - Hematopoietic stem cell transplantation is the only curative treatment; high mortality rate 	<ul style="list-style-type: none"> - Elevated WBCs (leukocytosis) - Genetic testing to identify mutations - Flow cytometry analysis to evaluate neutrophil expressions - Key clinical finding: absence of pus at site of infection 	<ul style="list-style-type: none"> - Aggressive and severe gingivitis and rapidly progressive periodontitis - Persistent oral ulcers (gingivostomatitis) - Absence of pus - Premature exfoliation of primary dentition and early loss of permanent teeth 	<p>Alveolar bone loss</p>	<ul style="list-style-type: none"> - Periodontal disease may be refractory to nonsurgical periodontal treatment and rigorous home care regimens - Prompt targeted antibiotic therapy - Adjunctive treatment may include granulocyte/thrombocyte transfusions, recombinant factor VIIa, and intravenous immunoglobulins - Prophylactic antibiotics prior to dental procedures
Obesity¹⁰	<ul style="list-style-type: none"> - Chronic complex multifactorial metabolic disorder presenting as excessive accumulation of fat - Etiologies: genetic, neuro-endocrine, drug-induced, behavioral (diet and activity) - Comorbidities: diabetes, hypertension, CV disease, obstructive sleep apnea, systemic inflammation, some cancers 	<ul style="list-style-type: none"> - Measured BMI <ul style="list-style-type: none"> • Ages 2-19: \geq 95th percentile or \geq 30 kg/m², whichever is lower based on age and gender using CDC growth charts • Adult: \geq 30 kg/m² - Other measures include: waist/hip circumferences, waist to hip ratios 	<ul style="list-style-type: none"> - Increased plaque index, bleeding on probing, periodontal pocket depth, clinical attachment loss - Mouth breathing 	<p>Alveolar bone loss</p>	<ul style="list-style-type: none"> - Comorbidities may influence management - Dietary weight loss may reduce systemic inflammation and, in turn, enhance response to periodontal therapy

Abbreviations in table: BMI: Body mass index; CDC: Centers for Disease Control and Prevention; CV: cardiovascular; NSAIDs: nonsteroidal anti-inflammatory drugs; PLS: Papillon Lefèvre syndrome; SRP: Scaling and root planing; WBCs: White blood cells.

Disorder	General characteristics	Diagnostic criteria	Oral findings		Treatment considerations
			Clinical	Radiographic	
Papillon Lefèvre syndrome (PLS)¹¹	<ul style="list-style-type: none"> - Autosomal recessive disorder - Palmoplantar hyperkeratosis, nail dystrophy, pyogenic skin and other infections, intracranial calcification - Rapidly progressing periodontal manifestations beginning shortly after tooth eruption; occurs in primary and permanent dentitions - Etiology: alterations in the CTSC gene and, likely, neutrophil defects 	<ul style="list-style-type: none"> - Urinalysis for cathepsin C activity - Genetic testing for mutation of cathepsin C gene - Key clinical finding: periodontal degeneration 	<ul style="list-style-type: none"> - Rapidly advancing gingival inflammation and bleeding, deep periodontal pockets, gingival abscesses, periodontal destruction - Premature loss of all primary teeth by ages 4-5; loss of permanent teeth by age 16 - After tooth loss, gingiva returns to healthy state 	Generalized extensive alveolar bone loss with migration of teeth	<ul style="list-style-type: none"> - Treatment of oral manifestations depends on patient's age, psychological state, and tooth mobility - May include nonsurgical therapy (e.g., monthly SRP, systemic antibiotics) and/or extraction of hopeless teeth - Alveolar loss renders prosthetic rehabilitation challenging
Periodontal Ehlers-Danlos syndrome^{12,13} (Synonyms: Ehlers-Danlos syndrome VIII; pEDS)	<ul style="list-style-type: none"> - One of a group of hereditary connective tissue disorders; autosomal dominant - Characterized by varying features including tissue fragility with easy bruising, vascular complications, joint hypermobility and/or pain, pretibial discoloration/plaques, increased infection rate, hoarse voice - Predominant feature is severe early-onset periodontitis (mean age 14) 	<ul style="list-style-type: none"> - Clinical examination and molecular genetic testing (variant in the genes C1R and C1S which play a role in innate immune system) - Complete lack of gingival attachment is considered pathognomonic - Most children identified through family history 	<ul style="list-style-type: none"> - Severe gingival inflammation, loss of attached gingiva, and gingival thinning and recession - Rapid alveolar bone loss - Premature tooth loss 	Alveolar bone loss	<ul style="list-style-type: none"> - May include nonsurgical therapy (e.g., monthly SRP, systemic antibiotics) and/or extraction of hopeless teeth - Alveolar loss renders prosthetic rehabilitation challenging - Implants at high risk of peri-implantitis

Abbreviations in table: BMI: Body mass index; CDC: Centers for Disease Control and Prevention; CV: cardiovascular; NSAIDs: nonsteroidal anti-inflammatory drugs; PLS: Papillon Lefèvre syndrome; SRP: Scaling and root planing; WBCs: White blood cells.

References

1. Thumbigere Math V, Rebouças P, Giovani PA, et al. Periodontitis in Chédiak-Higashi syndrome: An altered immunoinflammatory response. *JDR Clin Trans Res* 2018;3(1):35-46. Available at: "https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5734460/". Accessed September 21, 2022.
2. National Organization for Rare Disorders (NORD). Chédiak Higashi Syndrome. Available at: "https://rarediseases.org/rare-diseases/chediak-higashi-syndrome/". Accessed September 1, 2022.
3. American Diabetes Association. Diabetes Overview. Understanding A1c. Diagnosis. Available at: "https://diabetes.org/diabetes/a1c/diagnosis". Accessed September 21, 2022.
4. Albandar JM, Susin C, Hughes FJ. Manifestations of systemic diseases and conditions that affect the periodontal attachment apparatus: Case definitions and diagnostic considerations. *J Periodontol* 2018;89(Suppl 1):S183-S203.
5. National Organization for Rare Disorders (NORD). Haim-Munk Syndrome. Available at: "https://rarediseases.org/rare-diseases/haim-munk-syndrome/". Accessed September 1, 2022.
6. National Organization for Rare Disorders (NORD). Hypophosphatasia. Available at: "https://rarediseases.org/rare-diseases/hypophosphatasia/". Accessed September 1, 2022.
7. Okawa R, Kokomoto K, Nakano K. Dental effects of enzyme replacement therapy in case of childhood-type hypophosphatasia. *BMC Oral Health* 2021;21(1):323.
8. National Organization for Rare Disorders (NORD). Langerhans Cell Histiocytosis. Available at: "https://rarediseases.org/rare-diseases/papillon-lefevre-syndrome/". Accessed September 1, 2022.
9. National Organization for Rare Disorders (NORD). Leukocyte Adhesion Deficiency Syndromes. Available at: "https://rarediseases.org/rare-diseases/leukocyte-adhesion-deficiency-syndromes/". Accessed September 1, 2022.
10. National Collaborative on Childhood Obesity Research. A Guide to Methods for Assessing Childhood Obesity: Overview of Body Composition and Measuring Adiposity. Available at: "https://www.nccor.org/tools-assessingobesity-guide/overview-of-body-composition-and-measuring-adiposity/#two-c". Accessed August 24, 2022.
11. National Organization for Rare Disorders (NORD). Papillon Lefèvre Syndrome. Available at: "https://rarediseases.org/rare-diseases/papillon-lefevre-syndrome/". Accessed August 24, 2022.
12. Kapferer-Seebacher I, van Dijk FS, Zschocke J. Periodontal Ehlers-Danlos Syndrome. In: Adam MP, Everman DB, Mirzaa GM, et al., eds. *GeneReviews*[®] [Internet]. Seattle, Wash.: University of Washington, Seattle; 2021:1993-2022. Available at: "https://www.ncbi.nlm.nih.gov/books/NBK572429/#". Accessed August 24, 2022.
13. Rinner A, Zschocke J, Schossig A, Gröbner R, Strobl H, Kapferer-Seebacher I. High risk of peri-implant disease in periodontal Ehlers-Danlos Syndrome. A case series. *Clin Oral Implants Res* 2018;29:1101-6.